

## Expression alterations of KRAS gene associated with AgNPs

**Mohammed Ayyed Najm** 

\*Corresponding author: Assistant Professor (PhD), Department of Pharmacology, College of Medicine, Ibn Sina University of Medical and Pharmaceutical Sciences, Baghdad, Iraq. E-mail address: dr.mohammed.ayyed@ibnsina.edu.iq

**Gulboy Abdolmajeed Nasir** 

Assistant Professor (Ph.D.), College of Agricultural Engineering Sciences, University of Baghdad, Iraq. E-mail address: gulboy.nasir@coagri.uobaghdad.edu.iq

**Huda Musleh Mahmood** 

\*Corresponding author: Assistant Professor (Ph.D.) at Department of Biotechnology, College of Sciences, University of Anbar, Iraq. E-mail address: huda.mahmood@uoanbar.edu.iq

---

### ***Abstract***

#### **Objective**

Nanotechnology has had a significant impact on medicine, cosmetics, and food packaging via the utilization of artificially created nanomaterials like AgNPs. Due to their potent antibacterial properties, AgNPs are popular nanoparticles in consumer and medical products. This investigation sought to explore the effects of AgNPs on the expression of the Kirsten rat sarcoma (KRAS) viral gene in the liver and spleen of mice to evaluate the molecular safety of exposure to AgNPs.

#### **Materials and Methods**

A total of 56 mice were assigned to seven different groups (8 mice in each). Three groups were treated with AgNPs every day at 0.25, 0.5, and 1 mg per kg for one week. Remaining three groups were given the same amount of treatment for 14 days. One group functioned as an untreated counterpart. After the treatment, the spleen and liver tissues were excised. Later, total RNA was isolated using the method of Junqueira and Carneiro, and cDNA was made using the Applied Biosystem Kit (Product No. 4387406) for the measurement of gene expression in quantitative form.

## Results

The expression of the KRAS gene changed with the amount and length of time spent in AgNP. The expression of KRAS was decreased in all of the AgNP-treated in comparison to the control liver's tissue. On the other hand, the KRAS expression in the spleen of the treated animals was consistently higher than that of the controls. These modifications were more significant at higher concentrations and longer durations.

## Conclusions

The results demonstrated that AgNP publicity altered the expression of KRAS genes in a tissue-unique way: downregulation was observed inside the liver, and upregulation changed into found inside the spleen. These distinctive classes of expression suggest that AgNPs can influence cell communication pathways depending on the features of the organ and the microenvironment. Specifically, the sustained growth in KRAS in the spleen following the injection of AgNPs, particularly at higher doses and with a longer period of publicity, has caused large concerns regarding the ability of the injection to lead to a cancerous manner in related tissues, which are immune-associated. These findings display the need for careful consideration of AgNP's utilization, specifically in merchandise meant for long-term or worldwide use. Future planned investigations have to discover the capacity mechanisms through which AgNPs affect KRAS, assess the capability of long-term consequences of pathology, and determine the safe limits of exposure. Comprehensive molecular safety evaluation is crucial to ensuring the responsible development and usage of silver-primarily based technologies in scientific and commercial enterprise settings.

**Keywords:**  $\beta$ -ACTIN, gene expression, liver, silver nanoparticles, spleen

**Paper Type:** Research Paper.

**Citation:** Najm, M. A., Nasir, G. A., & Mahmood, H. M. (2025). Expression alterations of KRAS gene associated with AgNPs. *Agricultural Biotechnology Journal* 17(2), 217-236.

*Agricultural Biotechnology Journal* 17 (2), 217-236.

DOI: 10.22103/jab.2025.24920.1672

Received: March 02, 2025.

Received in revised form: April 20, 2025.

Accepted: April 22, 2025.

Published online: June 30, 2025.

Publisher: Faculty of Agriculture and Technology Institute of Plant Production, Shahid Bahonar University of Kerman-Iranian Biotechnology Society.



© the authors

## Introduction

The manufacturing and utilization of nanoparticles have enhanced appreciably over recent years, in general due to two motives: simple manufacturing of nanoparticles and their increasing range of applications (Al-Tae, 2020; Naief et al., 2023). Specifically, silver nanoparticles have come to be more popular most of the numerous forms of nanoparticles. Their sizeable reputation is attributed to their multiple homes, which include broad-spectrum antimicrobial interest in opposition to many microbial species (Tiàn et al., 2018; Kūmar et al., 2018; Kadhūm and Hussein, 2020), the incredibly low cost in their synthesis, and the flexibility to produce exceptionally sized and fashioned marketers (Sohn et al., 2015; Rajapantulu and Bandyopadhyaya, 2021). Today, AgNPs are not unusual in lots of everyday merchandise like textiles, meals containers, wound dressings, and personal care products (Wei et al., 2015; McGillicuddy et al., 2017). However, despite the numerous advantages of AgNPs, worries about their safety, mostly due to the capability of releasing ( $\text{Ag}^+$ ), which might be poisonous to organisms (Yue et al., 2017; Padhye et al., 2023). Despite the not unusual fact that the  $\text{Ag}^+$  ions release is accountable for the toxicity of AgNPs, some studies have observed a link among the two (Sorensen et al., 2015; Mat Lazim et al., 2023). Other scientists have even stated that the toxicity of metal AgNPs is greater than that of the ions themselves (Sakamoto et al., 2015; Pakrashi et al., 2017; Abramenko et al., 2018). The complexity indicates that the organic results of AgNPs are complicated and require further molecular research. An interesting gene here is Kirsten's viral oncogene for rats (KRAS), this gene is a proto-oncogene that encodes a small GTPase which takes part in the cell proliferation, differentiation, and survival. Alterations of KRAS are found in many human cancers; hence, it is an important target for research and development in oncology (Cox, 2014). Overexpression or mutation of KRAS has been associated with tumorigenesis, metastasis, and angiogenesis in cancer (Mahmood *et al.*, 2019; Gökmen *et al.*, 2023). Therefore, expression studies on environmental influences such as nanoparticles on KRAS provide implications both for toxicology as well as therapy. AgNPs showed anti-cancer activities, which involved down-regulation of cancer genes like KRAS. For example, reverse transcriptase-qPCR has proven that AgNPs substantially diminished the KRAS expression in treated cancer cells (Bin Saeed et al., 2023). On the other hand, increasing worries nearby the disadvantages of doses or long periods of exposure to chemical synthesis AgNPs, which can lead to cell toxicity and loss of efficacy as a treatment option (Srisaisap & Boonserm, 2024). As a result, while performing future studies in molecular terms for safety applications against cancer therapy applications, epigenetic modifications should also be taken into consideration. NPs can carry out gene expression modification through several recent studies on epigenetic mechanisms concerning DNA methylation and histone tail modifications, along with chromatin restructuring and noncoding RNAs like microRNAs and

long noncoding RNAs. The control of epigenetics is very complex, and the factors that contribute to it are diets, infections, climate conditions, and even xenobiotics (Amiri Roudbar et al., 2020). Such interactions occur on multiple levels and significantly define the expression profiles at genes that modulate phenotype and disease susceptibility (Safâei et al., 2022; Alavi et al., 2022). Gene regulation in eukaryotes managed tissue-specifically as well as time-specifically. At any given type of tissue, only a limited number of actively expressed genes; their expression pattern varies with developmental stage and health status (Hajalizadeh et al., 2021; Heidarpour et al., 2011). Both intrinsic factors, like chromatin architecture, as well as extrinsic elements such as inter-tissue signals, participate in this spatial and temporal control (Shokri et al., 2023). Epigenetic modifications serve as a mechanism through which environmental factors can trigger inheritable modifications in gene expression without changing the DNA sequence. Studies carried out reveal that maternal nutritional status during gestation has an imprinting effect on offspring epigenetics, at least regarding development, metabolism, and disease predisposition. Exposure of organisms to environmental stressors such as toxins and pathogens is also capable of inducing epigenetic modifications that affect gene expression and cellular behavior. This learning holds great value. Knowledge of epigenetics may improve livestock breeding, enhance productivity, and improve disease resistance (increased health resilience). Epigenetic marks are in use for forecasting health results and performance traits in diverse species (livestock included). The basic genetic research has now shifted towards the gene and protein identification related to specific traits, and then studying these genes and proteins via their chromosomal and cellular analysis (Bordbar et al., 2022). Increasing knowledge about the regulation of genes plus their response to environmental factors holds great promise for new therapeutic innovations that would enhance health in animals as well as humans (Arabpour et al., 2021; Mohammadabadi et al., 2022b; Louay et al., 2020). Therefore, this paper attempted a molecular safety evaluation of silver nanoparticles by investigating the impact of such particles on KRAS expression in mouse liver and spleen. The tissues were selected based on previously reported sensitivity to nanoparticle assays, along with their primary role in detoxification as well as immune regulation. This study tries to know the exact molecular impacts of AgNPs by looking at how different levels and times of exposure to AgNPs affect the KRAS gene expression in the various tissues; this will aid in giving better info about their safety.

## Material and methods

**AgNPs preparation:** The silver nanoparticles (AgNPs) used in this research have been previously utilized in the preceding observe by using Nasir et al. (2016) and described in detail.

Spherical AgNPs with an ordinary length of  $40 \pm 5$  nm were created using an environmentally friendly technique that utilized the extract of olives as the reducing agent and D-sorbitol as the stabilizer (Nasir et al., 2016; Nasir et al., 2020). The shape, length distribution, and balance of the synthesized nanoparticles were evaluated, and they were stored in a dark, wrapped box at  $37^{\circ}\text{C}$  till they were used in this experiment to prevent the formation of clusters and photodegradation.

**Animals used in experiments:** Adult male mice (weight variety: 24-36 g; age: 8-10 weeks) were involved in this study. The mice were housed in a monitored condition with a temperature. Of  $23\text{--}25^{\circ}\text{C}$  with a 12 hr. mild-dark cycle (Balls, 2022). The mice have been given a week to adapt to the laboratory's conditions before the experiments started (Klimek & Rogalska, 2021). A total of 56 mice had been assigned to 7 agencies of 8 mice each, these businesses were housed in separate cages. All mice were permitted to eat the standard pelleted feed and drink water freely at some point during the experimental period. The businesses that obtained remedies had been segregated into:

Group 1: 0.250 mg of AgNP / 7 days, Group 2: 0.500 mg of AgNP / 7 days, Group 3: 1,000 mg of AgNP / 7 days, Group 4: 0.250 mg of AgNP / 14 days, Group 5: 0.500 mg of AgNP / 14 days, Group 6: 1,000 mg of AgNP / 14 days, Group 7 (control group): 0.500 mg of distilled water for 14 days.

**Tissue sampling:** All mice were sacrificed 24 hours after the completion of their respective treatments using cervical dislocation as the euthanasia method (Balls, 2022). Hepatic and splenic tissues were immediately harvested for downstream molecular and histopathological analyses.

**RNA extraction and histological processing:** Tissue samples from the spleen and liver were processed for histopathological analysis following the approach described by Junqueira and Carneiro (Mescher, 2024). Tissue fragments were cut into small to medium-sized pieces with a thickness of 5 to 10  $\mu\text{m}$  using a precise microscope that rotates.

**cDNA synthesis:** Whole RNA was isolated from the tissues (liver and spleen), and cDNA using the Applied Biosystems High-Capacity cDNA Reverse Transcription Kit (Cat.No.4387406). The reactions were conducted in a thermal cycler according to the protocol outlined by the manufacturer, maintaining the temperature and reaction time as described in Table 1. The cDNA that was synthesized was kept at  $-20^{\circ}\text{C}$  for analysis of the gene expression.

**Primer design and qPCR analysis:** Primers for the intended gene (KRAS) and a housekeeping gene ( $\beta$ -ACTIN) were created using the Primer3Plus software (Table 2).  $\beta$ -ACTIN was chosen as the housekeeping gene to equalize the amount of the target gene (Li et al., 2010). The comparative Ct method ( $\Delta\Delta\text{Ct}$ ) was employed to assess the expression ratio of the target

gene. This approach contrasts the Ct values of the treated samples with those of the control samples and normalizes the latter to the internal reference gene  $\beta$ -ACTIN (Ali et al., 2022).


**Table 1. Reverse transcription thermal profile**

Temperature (°C)	Heated Lid	S-1	S-2	S-3	S-4	S-5
Time (min.)	—	10	10	60	5	$\infty$
Value	111	25	37	42	75	4

**Table 2. Primer sequences and melting temperatures for KRAS and  $\beta$ -ACTIN**

Gene	Primer	Sequence (5' → 3')	Tm (°C)
KRAS	Forward	AGGCCTGCTGAAAATGACTG	63.9
KRAS	Reverse	TCTGAATTAGCTGTATCGTCAAGG	65.9
$\beta$ -ACTIN	Forward	CCTGAACTACCTTGCAGCTC	60.0
$\beta$ -ACTIN	Reverse	ACTCCTGCTTGCTGATCCAC	63.0

**qPCR output:** Gene expression was evaluated through quantitative PCR, and results were visualized using amplification plots. Figure 1 illustrates the threshold cycle (Ct) values for each sample across all groups.



**Figure 1. CT values of KRAS and  $\beta$ -ACTIN gene expression across experimental groups**

## Results and discussion

The results of this investigation demonstrated that the KRAS gene expression in the liver and spleen increased significantly as a function of the concentration and duration of AgNP treatment. Also, the treatment group and the control group had different results (Table 3).

**Table 3.** The effects of one and two weeks of exposure to AgNPs in the spleen and liver tissues

Dose of AgNPs	Liver (Mean $\pm$ SE)	Spleen (Mean $\pm$ SE)
0.25 mg/kg bw / 1 week	0.01268 $\pm$ 0.0045 <sup>bc</sup>	0.0107 $\pm$ 0.0006 <sup>bc</sup>
0.5 mg/kg bw / 1 week	0.0118 $\pm$ 0.0030 <sup>bc</sup>	0.0254 $\pm$ 0.0041 <sup>ab</sup>
1.0 mg/kg bw / 1 week	0.0114 $\pm$ 0.0043 <sup>bc</sup>	0.0306 $\pm$ 0.0070 <sup>a</sup>
0.25 mg/kg bw / 2 weeks	0.0272 $\pm$ 0.0062 <sup>a</sup>	0.0172 $\pm$ 0.0087 <sup>abc</sup>
0.5 mg/kg bw / 2 weeks	0.0083 $\pm$ 0.0045 <sup>bc</sup>	0.0079 $\pm$ 0.0032 <sup>c</sup>
1.0 mg/kg bw / 2 weeks	0.0024 $\pm$ 0.0005 <sup>c</sup>	0.0068 $\pm$ 0.0051 <sup>c</sup>
Control	0.0145 $\pm$ 0.0022 <sup>b</sup>	0.0099 $\pm$ 0.0016 <sup>bc</sup>
<b>LSD</b>	<b>0.0117 *</b>	<b>0.0159 *</b>

\* Significant at  $P < 0.05$ . Different superscripts in each row designate significant differences between columns.

Administration of the lowest AgNP dose (0.25 mg/kg) for two weeks led to the highest hepatic KRAS gene expression value (0.0272), while the lowest value (0.0024) was recorded in the group that received the highest dose (1 mg/kg) for two weeks. This finding suggests that prolonged high-dose exposure may suppress KRAS gene expression in liver tissue, potentially indicating toxic or regulatory suppression mechanisms (Figure 2). In splenic tissue, the highest KRAS gene expression (0.0306) was observed following one-week exposure to 1 mg/kg AgNPs. On the contrary, the lowest splenic expression level (0.0068) was noted in the group that received the same dose for two weeks, reflecting a time-dependent decline despite high initial upregulation (Figure 3). Increasing proof supports the molecular toxicity of silver-based nanoparticles at low concentrations (Shen et al., 2015). Their outcomes on overall fitness, especially concerning the immune system, are documented (Shi et al., 2024). Our results confirmed a giant association between AgNP publicity and the expression of KRAS genes in tissue, particularly within the liver. Notably, the expression level of the KRAS gene inside the livers of mice that had been exposed

to one mg/kg AgNP for one week and 0.5-1 mg/kg AgNP for two weeks became substantially decrease.




Figure 2. Effect of AgNP treatments on hepatic KRAS gene expression




Figure 3. Effect of silver nanoparticle (AgNP) treatments on spleen KRAS gene expression in exclusive observation

Hepatocellular carcinoma (HCC) is the maximum not common shape of number one liver cancer and represents approximately 90% of all tumors inside the liver (Djuraeva et al., 2025). This malignancy is often related to the contamination of HBV and/or genetic adjustments in the RAS gene circle of relatives, such as mutations the Arzumanyan et al., 2013; Najm et al., 2020; Kouroumalis et al., 2023). KRAS mutations are in particular common in the liver's RAS family, they're not common than HRAS and NRAS mutations in this family of genes. Approximately 7% of all cancers have this type of mutation (Camonis et al., 2024). Previous investigations concur



with our findings. For instance, Bin Saeed et al. (2023) documented that AgNPs decreased the KRAS oncogene expression in most cancerous cells. Similarly, Xu et al. (2024) documented that silver nanoparticles had a sizable effect on lowering KRAS ranges. Elechalawar's team (2023) confirmed that nanoparticles can inhibit the activity of KRAS, which consequences in the suppression of tumor growth in cancers that have an imperative function of KRAS activation. In this research, the degrees of KRAS inside the liver tissues of the three experimental organizations were much lower than the control organization. Additionally, the average expression degree of KRAS in the three different-week agencies becomes constantly lower than the manipulate organization. These findings support the inhibitory role of AgNPs on the expression of KRAS inside the liver, this effect is mainly mentioned while the particles are uncovered for a long time.

On the other hand, the expression of KRAS within the spleen turned into associated with the AgNP dose throughout the one-week publicity duration. The KRAS gene expression tiers within the spleen of mice were zero.25 mg/kg AgNP for one week, and people dealt with 0.5-1 mg/kg AgNP for 2 weeks have been lower than those of the managed institution. However, a trendy in KRAS expression changed into observed in the one-week publicity group, which suggests that the duration of publicity has a great impact on the regulation of genes. KRAS is a sizable oncoprotein that participates in controlling the proliferation, transformation, and cellular specialization via various internalized pathways (Ma et al., 2019; Najm, 2019). Its improved expression is related to improved tumor growth and endothelial mobile development (Zhou et al., 2024; Nasir et al., 2025). As a result, the improved stage of KRAS located inside the spleen of mice handled with AgNPs for a restrained time period has brought about issues approximately its ability to sell cancer in other related organs. Despite the shortage of direct evidence that hyperlinks AgNP-prompted KRAS over expression to splenic most cancers, several studies have confirmed that AgNPs can reason great splenic harm. Rubab et al. (2023) found big changes in hematological tendencies, oxidative pressure signs, and microanatomy in rats that have been exposed to greenish artificial AgNPs.

Histological analysis demonstrated the presence of hemorrhage, vascular congestion, and an accumulation of brownish color in the purple pulp. These pigment deposits had been notably absent in the manipulated institution, which indicates that the spleen tissue is suffering from AgNP. The facts also tested that the expression stage of the KRAS gene in the spleen of the 3 one-week observe groups changed into higher than that of the manage organization, at the same time as the average expression level of the KRAS in spleen tissue for the 3 two-week observe groups turned into still beneath the manage organization's degree. This temporal shift may also endorse a strain response that results in gene upregulation, observed via a suppression or fatigue mechanism as the length increases.

Additionally, the position of the tumor's microenvironment in the disease process, particularly via genetic and epigenetic modifications, is documented. KRAS is mostly responsible for those changes, specifically in most cancers, which is related to irritation (Hazim et al., 2020; Raheem et al., 2021; Nasir et al., 2022; Abbas AL-Essawi and Mahmood, 2024; Daghaman et al., 2024). As KRAS upregulation promotes the improvement of tumors and promotes angiogenesis, our findings support the concept that exposure to AgNPs might also have a role in the early onset of cancerous behavior within the spleen tissue. Despite the lack of specific evidence of a hyperlink between AgNPs and splenic cancer, the observed changes in gene expression and spleen histology have to be studied in extra detail.

**Conclusions:** This research shows that the expression of KRAS is specifically altered in tissues following exposure to silver particles. Notably, the expression of KRAS in the liver was decreased following increased doses and longer exposures, which suggests that the molecule has a toxic or inhibitory effect. Conversely, a high-dose exposure period of short duration led to an increase in KRAS expression in the spleen, this was of concern because of the potential for it to lead to an immunological or carcinogenic outcome. These results concur with the theory that AgNPs have molecular effects that detrimentally affect the tissue and its function, as well as a stable genome. Future investigations should assess the long-term effects of these alterations and explain the molecular mechanisms behind the AgNP-induced alteration of gene expression.

### Conflict of Interest

There is no disagreement in this study.

### Authors' Contributions

Gulboy A. Nasir conducted the experiments in the laboratory and recorded the results. Mohammed A. Najm and Huda M. Mahmood had a role in the data analysis, writing the manuscripts, and developing the concept. All authors commented on and approved the ultimate version of the article.

### Acknowledgments

The authors would appreciate the insight of all reviewers and would appreciate their feedback, which helped to improve the quality and clarity of the paper.

## Ethical Considerations

All ethical guidelines were strictly followed throughout the study. The authors affirm that no data fabrication, falsification, plagiarism, or research misconduct occurred. Ethical approval was granted by the appropriate institutional committee (Approval No. 102, dated March 24, 2024).

## Funding

This study was conducted without external financial support and was entirely self-funded by the authors.

## References

- Abbas AL-Essawi, I., & Mahmood, H. M. (2024). Effect of gold nanoparticles on *hmgA* gene expression of *Pseudomonas aeruginosa* isolates. *Journal of Nanostructures*, 14(4), 1029-1039. <https://doi.org/10.22052/JNS.2024.04.004>
- Abramenko, N. B., Demidova, T. B., Abkhalimov, E. V., Ershov, B. G., Krysanov, E. Y., & Kustov, L. M. (2018). Ecotoxicity of different-shaped silver nanoparticles: Case of zebrafish embryos. *Journal of Hazardous Materials*, 347, 89-94. <https://doi.org/10.1016/j.jhazmat.2017.12.060>
- Alavi, M., Mozafari, M. R., Ghaemi, S., Ashengroph, M., Hasanzadeh Davarani, F., & Mohammadabadi, M. (2022). Interaction of epigallocatechin gallate and quercetin with spike glycoprotein (S-glycoprotein) of SARS-CoV-2: In silico study. *Biomedicines*, 10(12), e3074. <https://doi.org/10.3390/biomedicines10123074>
- Ali, F., Khan, A., Muhammad, S. A., & Hassan, S. S. U. (2022). Quantitative Real-Time Analysis of Differentially Expressed Genes in Peripheral Blood Samples of Hypertension Patients. *Genes*, 13(2), 187. <https://doi.org/10.3390/genes13020187>
- Al-taee, E. H. (2020). Effect of Silver Nanoparticles Synthesized Using Leaves Extract of Olive on Histopathological and Cytogenetic Effects in Albino Mice. *Iraqi Journal of Agricultural Sciences*, 51(5), 1448-1457. <https://doi.org/10.36103/ijas.v51i5.1155>
- Amiri Roudbar, M., Mohammadabadi, M. R., Ayatollahi Mehrgardi, A., Abdollahi-Arpanahi, R., Momen, M., Morota, G., Brito Lopes, F., Gianola, D., & Rosa, G. J. M. (2020). Integration of single nucleotide variants and whole-genome DNA methylation profiles for classification of rheumatoid arthritis cases from controls. *Heredity*, 124(5), 658-674. <https://doi.org/10.1038/s41437-020-0301-4>
- Arabpour, Z., Mohammadabadi, M., & Khezri, A. (2021). The expression pattern of p32 gene in femur, humeral muscle, back muscle and back fat tissues of Kermani lambs. *Agricultural Biotechnology Journal*, 13(4), 183-200. <https://doi.org/10.22103/jab.2022.18782.1371>

- Arzumanyan, A., Reis, H. M., & Feitelson, M. A. (2013). Pathogenic mechanisms in HBV- and HCV-associated hepatocellular carcinoma. *Nature Reviews Cancer*, 13(2), 123–135. <https://doi.org/10.1038/nrc3449>
- Balls, M. (2022). Alternatives to laboratory animals: Trends in replacement and the Three Rs. *Alternatives to Laboratory Animals (ATLA)*, 50(1), 10–26. <https://doi.org/10.1177/02611929221082250>
- Bin Saeed, H. A., Daghestani, M. H., Ambreen, K., Khan, S., & Shahid, M. (2023). Low dose of green synthesized silver nanoparticles is sufficient to cause strong cytotoxicity via its cytotoxic efficiency and modulatory effects on the expression of PIK3CA and KRAS oncogenes in lung and cervical cancer cells. *Journal of Cluster Science*, 34, 2471–2485. <https://doi.org/10.1007/s10876-022-02395-3>
- Bordbar, F., Mohammadabadi, M., Jensen, J., Xu, L., Li, J., & Zhang, L. (2022). Identification of candidate genes regulating carcass depth and hind leg circumference in Simmental beef cattle using Illumina Bovine Beadchip and next-generation sequencing. *Animals*, 12(9), e1103. <https://doi.org/10.3390/ani12091103>.
- Camonis, J. H., Aushev, V. N., Zueva, E., & Zalcman, G. (2024). A review and perspective paper: Ras oncogene gets modest, from kingpin to mere henchman. *Cellular and Molecular Life Sciences*, 81(1), 412. <https://doi.org/10.1007/s00018-024-05449-z>
- Cox, A. D., Fesik, S. W., Kimmelman, A. C., Luo, J., & Der, C. J. (2014). Drugging the undruggable RAS: Mission possible? *Nature Reviews Drug Discovery*, 13(11), 828–851. <https://doi.org/10.1038/nrd4389>
- Daghman, B., Kaddar, N., Najm, M. A., & Barakat, H. (2024). Comparison of a double dose of paracetamol tablets 500 mg with one paracetamol tablet 1000 mg. *Research Journal of Pharmacy and Technology*, 17(10), 4671–4676. <https://doi.org/10.52711/0974-360X.2024.00720>
- Djuraeva, N. M., Ikramov, A. I., Abdukhalimova, Kh. V., Hursanova, D. Kh., & Kholmiraev, M. A. (2025). *Hepatocellular carcinoma complicating liver cirrhosis: A case report with emphasis on vascular involvement*. <https://doi.org/10.5281/zenodo.14957003>
- Elechalawar, C. K., Rao, G., Gulla, S. K., Patel, M. M., Frickenstein, A., Means, N., Roy, R. V., Tsiokas, L., Asfa, S., Panja, P., Rao, C., Wilhelm, S., Bhattacharya, R., & Mukherjee, P. (2023). Gold nanoparticles inhibit macropinocytosis by decreasing KRAS activation. *ACS Nano*, 17(10), 9326–9337. <https://doi.org/10.1021/acsnano.3c00920>
- Gökmen, İ., Taştekin, E., Demir, N., Özcan, E., Akgül, F., Hacıoğlu, M. B., Erdoğan, B., Topaloğlu, S., & Çiçin, İ. (2023). Molecular pattern and clinical implications of

- KRAS/NRAS and BRAF mutations in colorectal cancer. *Current Issues in Molecular Biology*, 45(10), 7803–7812. <https://doi.org/10.3390/cimb45100491>
- Hajalizadeh, Z., Dayani, O., Khezri, A., Tahmasbi, R., Mohammadabadi, M., Solodka, T., Kalashnyk, O., Afanasenko, V., & Babenko, O. (2021). Expression of calpastatin gene in Kermani sheep using real-time PCR. *Journal of Livestock Science and Technologies*, 9(2), 51-57. <https://doi.org/10.22103/jlst.2021.18165.1381>
- Hazim, O. H., Najm, M. A., & Ibraheem, Z. O. (2020). Genetic diversity of three medicinal plants of Apiaceous family. *Systematic Reviews in Pharmacy*, 12, 737–741. <https://www.semanticscholar.org/paper/Genetic-Diversity-of-Three-Medicinal-Plants-of-Hazim-Najm/5a826b322f251f827618b9829c075aede5454055>
- Heidarpour, F., Mohammadabadi, M. R., Zaidul, I. S. M., Maherani, B., & Saari, N. (2011). Use of prebiotics in oral delivery of bioactive compounds: A nanotechnology perspective. *Pharmazie*, 66(5), 319-324. <https://pubmed.ncbi.nlm.nih.gov/21699064/>
- Kadhun, M. M., & Hussein, N. N. (2020). Detection of the antimicrobial activity of silver nanoparticles biosynthesized by *Streptococcus pyogenes* bacteria. *Iraqi Journal of Agricultural Sciences*, 51(2), 500–507. <https://doi.org/10.36103/ijas.v51i2.976>
- Khabiri, A., Toroghi, R., Mohammadabadi, M., & Tabatabaeizadeh, S. E. (2023). Introduction of a Newcastle disease virus challenge strain (sub-genotype VII.1.1) isolated in Iran. *Veterinary Research Forum*, 14(4), 221. <https://doi.org/10.30466/vrf.2022.548152.3373>
- Klimek, A., & Rogalska, J. (2021). Extremely low-frequency magnetic field as a stress factor—Really detrimental? Insight into literature from the last decade. *Brain Sciences*, 11(2), 174. <https://doi.org/10.3390/brainsci11020174>
- Kouroumalis, E., Tsomidis, I., & Voumvouraki, A. (2023). Pathogenesis of hepatocellular carcinoma: The interplay of apoptosis and autophagy. *Biomedicines*, 11(4), 1166. <https://doi.org/10.3390/biomedicines11041166>
- Kumar, S. S. D., Houreld, N. N., Kroukamp, E. M., & Abrahamse, H. (2018). Cellular imaging and bactericidal mechanism of green-synthesized silver nanoparticles against human pathogenic bacteria. *Journal of Photochemistry and Photobiology B: Biology*, 178, 259–269. <https://doi.org/10.1016/j.jphotobiol.2017.11.001>
- Li, Z., Yang, L., Wang, J., Shi, W., Pawar, R., Liu, Y., Xu, C., Cong, W., Hu, Q., Lu, T., Xia, F., & Guo, W. (2010).  $\beta$ -Actin is a useful internal control for tissue-specific gene expression studies using quantitative real-time PCR in the half-smooth tongue sole *Cynoglossus semilaevis* challenged with LPS or *Vibrio anguillarum*. *Fish & Shellfish Immunology*, 29(1), 89–93. <https://doi.org/10.1016/j.fsi.2010.02.021>

- Louay M. Al-Ani, Maha R. Ghreeb, Huda M. Mahmood, Bilal J.M. Aldahham. (2020). Teratogenicity of Pyocyanin Pigment Isolated from Local *Pseudomonas aeruginosa* Isolates on Mice Neural Tube Defects (NTDs) and other Abnormalities. *Systematic Reviews in Pharmacy*, 11 (7), 600-604. [doi:10.31838/srp.2020.7.84](https://doi.org/10.31838/srp.2020.7.84)
- Ma, Q., Gu, L., Liao, S., Zheng, Y., Zhang, S., Cao, Y., & Wang, Y. (2019). NG25, a novel inhibitor of TAK1, suppresses KRAS-mutant colorectal cancer growth in vitro and in vivo. *Apoptosis*, 24(1–2), 83–94. <https://doi.org/10.1007/s10495-018-1498-z>
- Mahmood, A. S., Hamid, S., & Najm, M. A. (2019). Association of polymorphism GST1 gene and antioxidant status, and interleukin-17 of colorectal cancer Iraqi patients. *Indian Journal of Forensic Medicine & Toxicology*, 13, 535. <https://www.researchgate.net/publication/337602852>
- Mat Lazim, Z., Salmiati, S., Marpongahtun, M., Arman, N. Z., Mohd Haniffah, M. R., Azman, S., Yong, E. L., & Salim, M. R. (2023). Distribution of silver (Ag) and silver nanoparticles (AgNPs) in aquatic environment. *Water*, 15(7), 1349. <https://doi.org/10.3390/w15071349>
- McGillicuddy, E., Murray, I., Kavanagh, S., Morrison, L., Fogarty, A., Cormican, M., Dockery, P., Prendergast, M., Rowan, N., & Morris, D. (2017). Silver nanoparticles in the environment: Sources, detection and ecotoxicology. *Science of the Total Environment*, 575, 231–246. <https://doi.org/10.1016/j.scitotenv.2016.10.041>
- Mescher, A. L. (Ed.). (2024). *Junqueira's basic histology: Text and atlas* (17th ed.). McGraw-Hill. <https://accessmedicine.mhmedical.com/content.aspx?bookid=3390&sectionid=281539243>
- Mohamadinejad, F., Mohammadabadi, M., Roudbari, Z., Eskandarynasab Siahkouhi, S., Babenko, O., Klopenko, N., Borshch, O., Starostenko, I., Kalashnyk, O., & Assadi Soumeh, E. (2024). Analysis of liver transcriptome data to identify the genes affecting lipid metabolism during the embryonic and hatching periods in ROSS breeder broilers. *Journal of Livestock Science and Technologies*, 12(2), 61-67. <https://doi.org/10.22103/jlst.2024.23814.1554>
- Mohammadabadi, M., & Asadollahpour Nanaei, H. (2021). Leptin gene expression in Raini Cashmere goat using Real Time PCR. *Agricultural Biotechnology Journal*, 13(1), 197-214. <https://doi.org/10.22103/jab.2021.17334.1305>
- Mohammadabadi, M., Babenko, I., Borshch, O., Kalashnyk, O., Ievstafieva, Y., & Buchkovska, V. (2024). Measuring the relative expression pattern of the *UCP2* gene in different tissues

- of the Raini Cashmere goat. *Agricultural Biotechnology Journal*, 16(3), 317-332.  
<https://doi.org/10.22103/jab.2024.24337.1627>
- Mohammadabadi, M., Golkar, A., & Askari Hesni, M. (2023). The effect of fennel (*Foeniculum vulgare*) on insulin-like growth factor 1 gene expression in the rumen tissue of Kermani sheep. *Agricultural Biotechnology Journal*, 15(4), 239-256.  
<https://doi.org/10.22103/jab.2023.22647.1530>
- Mohammadabadi, M., Kheyroodin, H., Latifi, A., & Babenko Ivanivna, O. (2022a). mRNA expression profile of DNAH1 gene in testis tissue of Raini Cashmere goat. *Agricultural Biotechnology Journal*, 14(3), 243-256. <https://doi.org/10.22103/jab.2022.20199.1428>
- Mohammadabadi, M., Shaban Jorjandy, D., Arabpoor Raghavadi, Z., Abareghi, F., Sasan, H. A., & Bordbar, F. (2022b). The role of fennel on DLK1 gene expression in sheep heart tissue. *Agricultural Biotechnology Journal*, 14(2), 155-170.  
<https://doi.org/10.22103/jab.2022.19402.1399>
- Najm, M. A. (2019). Evaluation of antioxidant (GSH, CAT, SOD) and MDA in Iraqi women with breast cancer. *Journal of Global Pharma Technology*, 11(7), 107–110.  
<https://www.jgpt.co.in/index.php/jgpt/article/view/2597>
- Najm, M. A., Al-Hadeithi, Z. S. M., & Salih, A. T. A. (2020). Correlating schizophrenia with DRD3 Ser9Gly or HTR2 receptor gene variants by using the RFLP method. *Indian Journal of Forensic Medicine and Toxicology*, 14(3), 2038–2043.  
<https://doi.org/10.37506/ijfmt.v14i3.10727>
- Nasir, G. A., Mohammed, A. K., & Samir, H. F. (2016). Biosynthesis and characterization of silver nanoparticles using olive leaves extract and sorbitol. *Iraqi Journal of Biotechnology*, 15(1), 22–32. <https://jige.uobaghdad.edu.iq/index.php/IJB/article/view/167>
- Nasir, G. A., Najm, M. A., & Hussein, A. L. (2020). The effect of silver nanoparticles on BRAF gene expression. *Systematic Reviews in Pharmacy*, 11(10), 570–575.  
<https://www.sysrevpharm.org/abstract/the-effect-of-silver-nanoparticles-on-braf-gene-expression-66388.html>
- Nasir, G., Khudhair, I., Najm, M. A., & Mahmood, H. (2022). Nanotechnology at the molecular level. *Al-Rafidain Journal of Medical Sciences*, 3, 71–74.  
<https://ajms.iq/index.php/ALRAFIDAIN/article/view/88>
- Nasir G., Najm M., Mahmood H. (2025). Association Between Silver Nanoparticle Dose and Brain or Renal NF-κB Gene Expression. *J Nanostructures*, 15(1), 249-254.  
[Doi.10.22052/JNS.2025.01.024](https://doi.org/10.22052/JNS.2025.01.024)

- Noori, A. N., Behzadi, M. R. B., & Mohammadabadi, M. R. (2017). Expression pattern of Rheb gene in Jabal Barez Red goat. *The Indian Journal of Animal Sciences*, 87(11), 1375–1378. <https://doi.org/10.56093/ijans.v87i11.75890>
- Padhye, L. P., Jasemizad, T., Bolan, S., Tsyusko, O. V., Unrine, J. M., Biswal, B. K., Balasubramanian, R., Zhang, Y., Zhang, T., Zhao, J., Li, Y., Rinklebe, J., Wang, H., Siddique, K. H. M., & Bolan, N. (2023). Silver contamination and its toxicity and risk management in terrestrial and aquatic ecosystems. *Science of the Total Environment*, 871, 161926. <https://doi.org/10.1016/j.scitotenv.2023.161926>
- Pakrashi, S., Tan, C., & Wang, W. X. (2017). Bioaccumulation-based silver nanoparticle toxicity in *Daphnia magna* and maternal impacts. *Environmental Toxicology and Chemistry*, 36(12), 3359–3366. <https://doi.org/10.1002/etc.3917>
- Raheem, G. L. A., Nasir, G., Alshammari, M. M. M., & Ghasemian, A. (2021). Antibacterial and antibiofilm effects of bismuth nanoparticles produced by *Bacillus subtilis* against multidrug-resistant *Pseudomonas aeruginosa*. *Malaysian Journal of Biochemistry and Molecular Biology*, 3, 15–22. <https://repository.uobaghdad.edu.iq/articles/hxZ214sBVTCNdQwCbOHm>
- Rajapantulu, A., & Bandyopadhyaya, R. (2021). Formation of gold nanoparticles in water-in-oil microemulsions: Experiment, mechanism, and simulation. *Langmuir*, 37. <https://doi.org/10.1021/acs.langmuir.1c00084>
- Roudbar, M.A., Mohammadabadi, M., & Salmani, V. (2015). Epigenetics: A new challenge in animal breeding. *Genetics in the Third Millennium*, 12(4), 3900–3914.
- Rubab, M., Zain, A., Mubeen, B., Tariq, H., Malik, A., & Arshad, N. (2023). Green synthesized FM-AgNPs lead to alterations in hematology, oxidative stress biomarkers, and microanatomy of liver and spleen in rats. *Brazilian Archives of Biology and Technology*, 66. <https://doi.org/10.1590/1678-4324-2023220098>
- Safaei, S. M. H., Dadpasand, M., Mohammadabadi, M., Atashi, H., Stavetska, R., Klopenko, N., & Kalashnyk, O. (2022). An *Origanum majorana* leaf diet influences *Myogenin* gene expression, performance, and carcass characteristics in lambs. *Animals*, 13(1), 14. <https://doi.org/10.3390/ani13010014>
- Safaei, S. M. H., Mohammadabadi, M., Moradi, B., Kalashnyk, O., Klopenko, N., Babenko, O., Borshch, O. O., & Afanasenko, V. (2024). Role of fennel (*Foeniculum vulgare*) seed powder in increasing testosterone and *IGF1* gene expression in the testis of lamb. *Gene Expression*, 23(2), 98-105. <https://doi.org/10.14218/GE.2023.00020>



- Sakamoto, M., Ha, J. Y., Yoneshima, S., Kataoka, C., Tatsuta, H., & Kashiwada, S. (2015). Free silver ion is the main cause of acute and chronic toxicity of silver nanoparticles to cladocerans. *Archives of Environmental Contamination and Toxicology*, 68(3), 500–509. <https://doi.org/10.1007/s00244-014-0091-x>
- Shahsavari, M., Mohammadabadi, M., Khezri, A., Asadi Fozi, M., Babenko, O., Kalashnyk, O., Oleshko, V., & Tkachenko, S. (2023). Correlation between insulin-like growth factor 1 gene expression and fennel (*Foeniculum vulgare*) seed powder consumption in muscle of sheep. *Animal Biotechnology*, 34(4), 882-892. <https://doi.org/10.1080/10495398.2021.2000997>
- Shahsavari, M., Mohammadabadi, M., Khezri, A., Borshch, O., Babenko, O., Kalashnyk, O., Afanasenko, V., & Kondratiuk, V. (2022). Effect of fennel (*Foeniculum vulgare*) seed powder consumption on *insulin-like growth factor 1* gene expression in the liver tissue of growing lambs. *Gene Expression*, 21(2), 21-26. <https://doi.org/10.14218/GE.2022.00017>
- Shen, M. H., Zhou, X. X., Yang, X. Y., Chao, J. B., Liu, R., & Liu, J. F. (2015). Exposure medium: Key in identifying free Ag<sup>+</sup> as the exclusive species of silver nanoparticles with acute toxicity to *Daphnia magna*. *Scientific Reports*, 5, 9674. <https://doi.org/10.1038/srep09674>
- Shi, Q., Xue, C., Zeng, Y., Yuan, X., Chu, Q., Jiang, S., Wang, J., Zhang, Y., Zhu, D., & Li, L. (2024). Notch signaling pathway in cancer: From mechanistic insights to targeted therapies. *Signal Transduction and Targeted Therapy*, 9, 128. <https://doi.org/10.1038/s41392-024-01828-x>
- Shokri, S., Khezri, A., Mohammadabadi, M., & Kheyroodin, H. (2023). The expression of MYH7 gene in femur, humeral muscle and back muscle tissues of fattening lambs of the Kermani breed. *Agricultural Biotechnology Journal*, 15(2), 217-236. <https://doi.org/10.22103/jab.2023.21524.1486>
- Sohn, E. K., Johari, S. A., Kim, T. G., Kim, J. K., Kim, E., Lee, J. H., Chung, Y. S., & Yu, I. J. (2015). Aquatic toxicity comparison of silver nanoparticles and silver nanowires. *BioMed Research International*, 2015, 893049. <https://doi.org/10.1155/2015/893049>
- Sorensen, S. N., Hjorth, R., Delgado, C. G., Hartmann, N. B., Sørensen, S. N., & Baun, A. (2015). Nanoparticle ecotoxicity—Physical and/or chemical effects. *Integrated Environmental Assessment and Management*, 11, 722–724. <https://doi.org/10.1002/ieam.1683>
- Srisaisap, M., & Boonserm, P. (2024). Anticancer efficacy of biosynthesized silver nanoparticles loaded with recombinant truncated parasporin-2 protein. *Scientific Reports*, 14(1), 15544. <https://doi.org/10.1038/s41598-024-66650-5>
- Tian, X., Jiang, X., Welch, C., Croley, T. R., Wong, T. Y., Fan, S., Chong, Y., Li, R., Ge, C., Chen, C., & Yin, J. J. (2018). Bactericidal effects of silver nanoparticles on *Lactobacilli*

and the underlying mechanism. *ACS Applied Materials & Interfaces*, 10(10), 8443–8450.  
<https://doi.org/10.1021/acsami.7b17274>


Wei, L., Lu, J., Xu, H., Patel, A., Chen, Z. S., & Chen, G. (2015). Silver nanoparticles: Synthesis, properties, and therapeutic applications. *Drug Discovery Today*, 20(5), 595–601.  
<https://doi.org/10.1016/j.drudis.2014.11.014>

Xu, G., Song, X., Wang, X., Xue, R., Yan, X., Qin, L., Chang, X., Gao, J., Chen, Z., & Song, G. (2024). Combined miR-181a-5p and Ag nanoparticles are effective against oral cancer in a mouse model. *International Journal of Nanomedicine*, 19, 9227–9253.  
<https://doi.org/10.2147/IJN.S458484>

Yue, Y., Li, X., Sigg, L., Suter, M. J., Pillai, S., Yue, Y., Behra, R., & Schirmer, K. (2017). Interaction of silver nanoparticles with algae and fish cells: A side by side comparison. *Journal of Nanobiotechnology*, 15(1), 16. <https://doi.org/10.1186/s12951-017-0254-9>

Zhou, Y., Kuang, Y., Wang, C., Yu, Y., Pan, L., & Hu, X. (2024). Impact of KRAS mutation on the tumor microenvironment in colorectal cancer. *International Journal of Biological Sciences*, 20(5), 1947–1964. <https://doi.org/10.7150/ijbs.88779>

## تغییرات بیان ژن KRAS مرتبط با نانوذرات نقره (AgNPs)

محمد عاید نجم 


\*نویسنده مسئول: استادیار، گروه داروشناسی، دانشکده پزشکی، دانشگاه علوم پزشکی و داروسازی ابن سینا، بغداد، عراق. ایمیل:

dr.mohammed.ayyed@ibnsina.edu.iq

گلپوی عبدالمجید ناصر 

استادیار (دکتری)، دانشکده علوم مهندسی کشاورزی، دانشگاه بغداد، عراق. ایمیل:

gulboy.nasir@coagri.uobaghdad.edu.iq

هدی مصلح محمود 

\*نویسنده مسئول: استادیار (دکتری)، گروه بیوتکنولوژی، دانشکده علوم، دانشگاه الانبار، عراق. ایمیل:

huda.mahmood@uoanbar.edu.iq

تاریخ دریافت: ۱۴۰۳/۱۲/۱۲ تاریخ دریافت فایل اصلاح شده نهایی: ۱۴۰۴/۰۱/۳۱ تاریخ پذیرش: ۱۴۰۴/۰۲/۰۲

### چکیده

**هدف:** فناوری نانو انقلابی در حوزه‌های مختلف از جمله پزشکی، لوازم آرایشی و بسته‌بندی مواد غذایی ایجاد کرده است، به‌ویژه از طریق استفاده از نانومواد مهندسی‌شده مانند نانوذرات نقره (AgNPs). نانوذرات نقره به دلیل خواص ضد میکروبی قوی، از پرکاربردترین نانوذرات در محصولات مصرفی و زیست‌پزشکی محسوب می‌شوند. این مطالعه با هدف بررسی اثرات نانوذرات نقره بر سطح بیان ژن سرطان‌زای ویروسی Kirsten rat sarcoma (KRAS) در بافت‌های کبد و طحال موش‌ها انجام شد تا ایمنی مولکولی تماس با AgNPs ارزیابی شود.

**مواد و روش‌ها:** در مجموع ۵۶ موش به‌صورت تصادفی به هفت گروه (n=8 در هر گروه) تقسیم شدند. سه گروه به‌مدت ۷ روز مقادیر ۰/۲۵، ۰/۵ و ۱ میلی‌گرم بر کیلوگرم وزن بدن AgNP دریافت کردند. سه گروه دیگر همان دوزها را به‌مدت ۱۴ روز دریافت کردند. یک گروه نیز به عنوان کنترل بدون درمان باقی ماند. پس از پایان دوره درمان، بافت‌های کبد و طحال جمع‌آوری شد. RNA کل با استفاده از روش جوناکویرا و کارنیرو استخراج و سنتز cDNA با استفاده از کیت Applied Biosystem به شماره ۴۳۸۷۴۰۶ برای آنالیز بیان ژن انجام شد.

**نتایج:** سطح بیان ژن KRAS بسته به دوز و مدت زمان تماس با AgNP ها متفاوت بود. در بافت کبد، بیان ژن KRAS در همه گروه‌های دریافت‌کننده AgNP نسبت به گروه کنترل کاهش یافت. در مقابل، در بافت طحال، بیان ژن KRAS در تمام گروه‌های دریافت‌کننده به‌طور قابل توجهی افزایش داشت. این تغییرات در دوزهای بالاتر و مدت تماس طولانی‌تر شدیدتر بودند.

**نتیجه‌گیری:** یافته‌ها نشان می‌دهند که تماس با AgNP ها باعث تنظیم مجدد بیان ژن KRAS به‌صورت مختص-بافتی می‌شود؛ به‌طوری‌که کاهش بیان در بافت کبد و افزایش بیان در بافت طحال مشاهده شد. این الگوهای متفاوت بیان بیانگر آن هستند که نانوذرات نقره ممکن است مسیرهای پیام‌رسانی سلولی را به شیوه‌ای تحت تأثیر عملکرد اندام‌ها و ریزمحیط‌ها تغییر دهند. به‌ویژه، افزایش مداوم بیان KRAS در طحال پس از مصرف AgNP ها، به‌ویژه در دوزهای بالا و تماس طولانی‌مدت نگرانی‌هایی جدی درباره اختلالات ایمنی یا آغاز فرآیندهای سرطان‌زایی در بافت‌های مرتبط با سیستم ایمنی ایجاد می‌کند. این نتایج لزوم ارزیابی دقیق‌تر استفاده از AgNP ها را به‌ویژه در محصولاتی که برای کاربرد طولانی‌مدت یا سیستمیک طراحی شده‌اند، نشان می‌دهد. مطالعات بیشتری برای روشن‌سازی مکانیسم‌های زیربنایی تنظیم KRAS توسط AgNP ها، ارزیابی پیامدهای آسیب‌شناسی احتمالی بلندمدت، و تعیین آستانه‌های ایمن تماس ضروری است. تهیه یک پروفایل جامع ایمنی مولکولی برای تضمین توسعه و استفاده مسئولانه از فناوری‌های مبتنی بر نانوذرات نقره در حوزه‌های پزشکی و تجاری حیاتی است.

**کلمات کلیدی:** بیان ژن، طحال، کبد، نانوذرات نقره،  $\beta$ -ACTIN

**نوع مقاله:** پژوهشی.

**استناد:** محمد عاید نجم، گلبوی عبدالمجید ناصر، هدی مصلح محمود (۱۴۰۴). تغییرات بیان ژن KRAS مرتبط با نانوذرات نقره (AgNPs). مجله بیوتکنولوژی کشاورزی، ۱۷(۲)، ۲۱۷-۲۳۶.

Publisher: Faculty of Agriculture and Technology Institute of Plant Production, Shahid Bahonar University of Kerman-Iranian Biotechnology Society.



© the authors